

ACCEPTED

FOR REGISTRATION

April 03, 2013

New York State Department  
of Environmental Conservation  
Division of Materials Management  
Pesticide Product Registration

46001AL0022



EPA Est. No. 42750-MO-001

# KRENITE® S

## Brush Control Agent Water-Soluble Liquid

PEEL HERE TO OPEN



NET CONTENTS: 2.5 GALLONS

Manufactured for:

**ALBAUGH, INC.**  
1525 NE 36th Street  
Ankeny, Iowa 50021

**FOR CHEMICAL SPILL, LEAK,  
FIRE, OR EXPOSURE, CALL  
CHEMTREC (800) 424-9300**

ACTIVE INGREDIENT:	BY WEIGHT
Ammonium salt of fosamine [ethyl hydrogen (aminocarbonyl) phosphonate] . . . . .	41.5%
<b>OTHER INGREDIENTS</b> . . . . .	58.5%
<b>TOTAL</b> . . . . .	100.0%

Contains 4 Lbs. Active Ingredient per Gallon.

EPA Reg. No. 42750-247

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

**FIRST AID**

IF IN EYES:	
	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>

Have the product container label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for medical emergencies involving this product.

See inside booklet for additional PRECAUTIONARY STATEMENTS.

PF-65303

### PROOF

**THIS PROOF IS TO BE CHECKED FOR ACCURACY**

Please review and approve **Text, Spelling, Copy Placement, Size, Shape, Colors and Deline.**

**Authorized signature** accepts responsibility for accuracy of all copy, color break and artwork. Cimarron Label is not liable for any discrepancies subsequently identified.

**PLEASE NOTE:** Due to color variance between printers/monitors, the colors represented by this proof cannot be deemed accurate. Please refer to a color matching system such as the Pantone Matching System for a truer representation of spot colors.

**THIS PROOF IS NOT ACCURATE FOR COLOR-MATCH.**  
*Deline does not print.*

DATE	JOB NUMBER	CUSTOMER
January 18, 2013	65303	ALBAUGH, INC.
LABEL SIZE	LEAFLET FOLDED SIZE	LEAFLET FLAT SIZE
6.75" X 6.75"	6.5" X 6.0"	6.5" X 29.687"
LABEL COLORS	LEAFLET "OUT" COLORS	LEAFLET "IN" COLORS
PATTERN VARNISH: <input checked="" type="checkbox"/> YES <input type="checkbox"/> NO		

Form: CS 006L - 11/8/2011

**ARTWORK IS APPROVED**

**REVISED PROOF NEEDED**

**Cimarron Label**  
Experts in Extended Text Labeling

4201 North Westport Ave. • Sioux Falls, SD 57107  
Phone: (605) 978-0451 • Fax: (605) 978-0463

**WE CANNOT PROCESS  
THIS ORDER WITHOUT AN  
AUTHORIZED SIGNATURE**

Signed \_\_\_\_\_

Date \_\_\_\_\_

**Agri Star®**  
By Albaugh Inc.

# KRENITE® S

## Brush Control Agent

### Water-Soluble Liquid

Manufactured for:  
**ALBAUGH, INC.**  
1525 NE 36th Street  
Ankeny, Iowa 50021

**FOR CHEMICAL SPILL, LEAK,  
FIRE, OR EXPOSURE, CALL  
CHEMTREC (800) 424-9300**

AD052510  
PRODUCT OF CHINA

ACTIVE INGREDIENT:	BY WEIGHT
Ammonium salt of fosamine [ethyl hydrogen (aminocarbonyl) phosphonate]	41.5%
<b>OTHER INGREDIENTS</b>	<b>58.5%</b>
<b>TOTAL</b>	<b>100.0%</b>

Contains 4 Lbs. Active Ingredient per Gallon.

EPA Reg. No. 42750-247

## KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

### FIRST AID

IF IN EYES:	
	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>

Have the product container label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for medical emergencies involving this product.

**See inside booklet for additional PRECAUTIONARY STATEMENTS.**



---

## PRECAUTIONARY STATEMENTS

---

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION!** Causes (moderate) eye injury (irritation). Avoid contact with eyes or clothing.

---

### PERSONAL PROTECTIVE EQUIPMENT (PPE)

---

**Applicators and other handlers must wear:**

1. Long-sleeved shirt and long pants
2. Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

---

### USER SAFETY RECOMMENDATIONS

**USERS SHOULD:** Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.

---

### ENVIRONMENTAL HAZARDS

---

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters.

---

### PRODUCT INFORMATION

---

KRENITE® S brush control agent is a water-soluble liquid to be diluted with water and applied as a foliar spray for control and/or suppression of many woody species.

KRENITE® S may be applied for use in pine plantations and non-crop sites, including highway rights-of-way, industrial sites, railroad rights-of-way, storage areas, utility and pipeline rights-of-way.

This product may be applied in pine plantations and non-crop sites that contain areas of temporary surface water caused by collection of water between planting beds, in equipment ruts, or in other depressions created by management activities. It is permissible to treat intermittent drainage, intermittently flooded low-lying sites, seasonally dry flood plains and transitional areas between upland and lowland sites when no water is present. It is also permissible to treat marshes, swamps and bogs after water has receded, as well as seasonally dry flood deltas. **DO NOT** make applications to natural or man-made bodies of water, such as lakes, reservoirs, ponds, streams and canals.

KRENITE® S is non-flammable and nonvolatile.

---

### ENVIRONMENTAL CONDITIONS AND BIOLOGICAL ACTIVITY

---

A KRENITE® S spray directed to only part of susceptible brush species will provide control of the portion sprayed, resulting in a trimming effect. Treatment with KRENITE® S generally does not immediately affect deciduous woody plants; they retain normal foliage for the remainder of the growing season. Treated susceptible plants do not produce foliage or grow the following spring. Coniferous species treated with KRENITE® S generally displays visible symptoms following application.

Effectiveness may be reduced if, following treatment, rainfall occurs on the same day.

---

### RESISTANCE

---

When herbicides that affect the same biological site of action are used repeatedly over several years to control the same weed species in the same field, naturally-occurring resistant biotypes may survive a correctly applied herbicide treatment, propagate, and become dominant in that field. Adequate control of these resistant weed biotypes cannot be expected. If weed control is unsatisfactory, it may be necessary to retreat the problem area using a product affecting a different site of action.

To better manage herbicide resistance through delaying the proliferation and possible dominance of herbicide resistant weed biotypes, it may be necessary to change cultural practices within and between crop seasons such as using a combination of tillage, retreatment, tank-mix partners and/or sequential herbicide applications that have a different site of action. Weed escapes that are allowed to go to seed will promote the spread of resistant biotypes.

It is advisable to keep accurate records of pesticides applied to individual fields to help obtain information on the spread and dispersal of resistant biotypes. Consult your agricultural dealer, consultant, applicator, and/or appropriate state agricultural extension service representative for specific alternative cultural practices or herbicide recommendations available in your area.

---





---

## INTEGRATED PEST MANAGEMENT

---

This product may be used as part of an Integrated Pest Management (IPM) program that can include biological, cultural, and genetic practices aimed at preventing economic pest damage. IPM principles and practices include field scouting or other detection methods, correct target pest identification, population monitoring, and treating when target pest populations reach locally determined action thresholds. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action treatment threshold levels for treating specific pest/crop systems in your area.

---

## TANK MIXES

---

KRENITE® S herbicide may be tank mixed with other herbicides and/or adjuvant registered for use in pine plantations and non-crop sites. Follow all use directions, precautions, and restrictions on labels of tank-mixed products.

---

## SPRAY EQUIPMENT

---

KRENITE® S may be applied using high volume or low volume ground sprayers as well as aircraft (helicopter only). Application equipment must be calibrated before making applications of KRENITE® S.

---

## SPRAY ADJUVANTS

---

A penetrating type oil-based adjuvant (surfactant or crop oil concentrate) may be used with KRENITE® S. The adjuvant should be mixed in the spray solution at a minimum concentration of 1/4% by volume (1 quart per 100 gallons of spray solution) or at the manufacturer's recommended dosage.

If foaming is a problem during mixing, an anti-foam agent may be added.

---

## DIRECTIONS FOR USE

---

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Do not apply this product through any type of irrigation system.

Do not use on food or feed crops.

KRENITE® S must be used only in accordance with the labeling, or in supplemental Albaugh, Inc. labeling.

---

## AGRICULTURAL USES

---

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

1. Coveralls
2. Shoes plus socks

### PINE PLANTATIONS PREPARATION SITE

KRENITE® S may be applied for the post-harvest (pre-plant) control of undesirable pine and hardwood seedlings and saplings and suppression of brush and vines to aid site planting preparation for southern pines and/or genetically improved pines.





## APPLICATION INFORMATION

Apply as a foliar spray from mid-summer to when the target tree pests begin defoliation in late summer or fall. Applications of KRENITE® S may be made by ground or air (helicopter only) equipment. Use sufficient water to ensure complete coverage of the vegetation, 20 to 50 gallons per acre by ground and 10 to 15 gallons per acre by air.

## USE RATES AND PLANTS CONTROLLED

### Pine Seedlings and Saplings

Apply 2 to 4 quarts of KRENITE® S per acre for the control of seedling and sapling pines when burning is allowed on the site.

Apply 4 to 6 quarts per acre of KRENITE® S to control seedling and sapling pines when burning is not allowed on the site.

Use the higher rate when either pine saplings predominate or when high infestations of seedling pines are in the area to be sprayed.

### Combinations of Pine and Hardwood Seedlings and Saplings

To control a combination of pine and hardwood seedlings and saplings, apply a tank mixture of KRENITE® S at use rates indicated for spraying pine seedlings and saplings plus Imazapyr (4 pound active per gallon) at 8 to 20 ounces per acre. This tank mix may be applied for the control of Ash, Blackberry, Black gum, Black locust, Box elder, Cherry, Dogwood, Elms (winged, slippery), Oaks (red, white), Red maple, Sassafras, and Sourwood.

Follow all use directions, precautions and restrictions on Imazapyr product labels.

### Brush and Vine Suppression

The application of KRENITE® S plus Imazapyr will also provide suppression of brush and vines, such as, American beautyberry (French mulberry), Baccharis (groundsel tree), Vaccinium (blueberry) species, Wax myrtle (bayberry) and Wild grape.

\*Suppression – a visible reduction in plant population and/or plant vigor as compared to an untreated area and generally not accepted as control.

Do not apply more than 3 gallons of KRENITE® S per acre per year.

## NON-AGRICULTURAL USES

### NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow entry into treated areas until sprays have dried to perform hand tasks.

### NON-CROP SITES

KRENITE® S may be applied for general weed control as follows: uncultivated non-agricultural areas (such as airports, highway, railroad and utility rights-of-way, sewage disposal areas); uncultivated agricultural areas (non-crop producing, which includes: farmyards, fuel storage areas, fence rows, barrier strips); industrial sites (outdoor, such as lumberyards, pipeline and tank farms).

## APPLICATION INFORMATION

Make a foliar application of the recommended rate of KRENITE® S from full leaf expansion in the spring to the development of full canopy coloration in the fall for deciduous species to be controlled. Coniferous species, listed in the "USE RATES AND PLANTS CONTROLLED" chart below, may be treated at anytime during the growing season.

### LOW- AND HIGH-VOLUME DIRECTED SPRAYS

Prepare either a low-volume or high-volume spray solution of KRENITE® S. For the low-volume directed spray application, do not exceed a spray concentration of 30% by volume. For the high-volume directed spray application, do not use a spray concentration of less than 1.5% by volume.

Prepare the desired volume of spray solution by mixing the amount of this product in water as shown in the Spray Solution Table. Apply a quantity of spray solution which will thoroughly and uniformly cover the target plant foliage (spray to wet for high-volume applications). Rate and volume per acre will depend on the plant species, the height and density of plant growth as well as the type of application equipment used. On tall or dense stands of brush it may be necessary to spray from opposite sides in order to obtain thorough coverage of the foliage. Use the higher rate range on stands where difficult-to-control species are dominant. See the "USE RATES AND PLANTS CONTROLLED" section of the label for use rates and a listing of "difficult-to-control species."

Do not apply more than 6 gallons of KRENITE® S per acre per year.





### AERIAL and BROADCAST APPLICATIONS

Prepare a spray solution using 1-1/2 to 3 gallons of KRENITE® S in 10 to 40 gallons of water (see Spray Solution Table). For broadcast ground applications, use this product at the rate of 1.5 to 6 gallons per acre. Do not apply more than 6 gallons per acre when using ground equipment. For aerial applications, use this product at the rate of 1.5 to 3 gallons per acre. Do not apply more than 3 gallons of KRENITE® S per acre when using aerial equipment. Use sufficient spray volume to uniformly and thoroughly cover the foliage. Use the higher concentrations on stands in which difficult-to-control species are predominant (see "USE RATES AND PLANTS CONTROLLED" section for a listing of \*\*difficult-to-control species).

### SPRAY SOLUTION TABLE

Desired Volume	Amount of KRENITE® S						
	1.5%	2%	3%	4%	10%	20%	30%
5 Gal	**	**	**	0.8 qt	0.5 gal	1 gal	1.5 gal
10 Gal	0.6 qt	0.8 qt	1.2 qt	1.6 qt	1 gal	2 gal	3 gal
20 Gal	1.2 qt	1.6 qt	0.6 gal	0.8 gal	2 gal	4 gal	6 gal
30 Gal	0.45 gal	0.6 gal	0.9 gal	1.2 gal	3 gal	6 gal	**
40 Gal	0.6 gal	0.8 gal	1.2 gal	1.6 gal	4 gal	**	**
50 Gal	0.75 gal	1 gal	1.5 gal	2 gal	5 gal	**	**
100 Gal	1.5 gal	2 gal	3 gal	4 gal	**	**	**

### USE RATES AND PLANTS CONTROLLED

KRENITE® S effectively controls or suppresses (\*\*difficult-to-control listings) the following plants when applied at the use rates shown.

\*\*Suppression – a visible reduction in plant population and/or plant vigor as compared to an untreated area and generally not accepted as control.

1-1/2 to 6 gal KRENITE® S per acre			
Alder, red	<i>Alnus rubra</i>	Oak, red	<i>Quercus rubra</i>
Ash, white	<i>Fraxinus Americana</i>	Oak, water	<i>Quercus arkansana</i>
Aspen, quaking	<i>Populus tremuloides</i>	Oak, white	<i>Quercus alba</i>
Birch	<i>Betula</i> sp.	Persimmon**	<i>Diospyros virginiana</i>
Blackberry	<i>Rubus</i> sp.	Pine, loblolly	<i>Pinus taeda</i>
Blackgum	<i>Nyssa sylvatica</i>	Pine, Virginia	<i>Pinus virginiana</i>
Cherry, black**	<i>Prunus serotina</i>	Poplar, yellow (tulip tree)**	<i>Liriodendron tulipifera</i>
Cherry, pin	<i>Prunus pensylvanica</i>	Salmonberry	<i>Rubus spectabilis</i>
Chokecherry, common**	<i>Prunus virginiana</i>	Sassafras**	<i>Sassafras sassafras</i>
Elm**	<i>Ulmus</i> sp.	Sourwood**	<i>Oxydendrum arboretum</i>
Fern, bracken	<i>Pteridium aquilinum</i>	Spurge, leafy***	<i>Euphorbia esula</i>
Hawthorn**	<i>Crataegus</i> sp.	Sumac	<i>Rhus</i> sp.
Hickory**	<i>Carya</i> sp.	Sweetgum	<i>Liquidambar styraciflua</i>
Locust, black	<i>Robinia pseudoacacia</i>	Tallow, Chinese	<i>Sapium Sebiferum</i>
Maple, bigleaf**	<i>Acer macrophyllum</i>	Thimbleberry	<i>Rubus parviflorus</i>
Maple, red**	<i>Acer rubrum</i>	Willow**	<i>Salix</i> sp.
Maple, vine	<i>Acer circinatum</i>		
2 to 6 gal KRENITE® S per acre			
Basswood, American**	<i>Tilia Americana</i>	Grape, wild	<i>Vitis</i> sp.
Bindweed, field***	<i>Convolvulus arvensis</i>	Pine, Eastern white	<i>Pinus strobes</i>
Cottonwood, Eastern	<i>Populus deltoids</i>	Plum, wild	<i>Prunus munsoniana</i>
Elder, American	<i>Sambucus canadensis</i>	Rose, multiflora	<i>Rosa multiflora</i>
Elm, slippery	<i>Ulmus rubra</i>	Sycamore	<i>Platanus occidentalis</i>
Elm, winged**	<i>Ulmus alata</i>	Tree-of-heaven	<i>Ailanthus altissima</i>

\*\*Difficult-to-control or Suppression

Suppression – A visible reduction in plant population and/or plant vigor as compared to an untreated area and generally not accepted as control.

\*\*\*Make applications after plants begin to bloom.



## TANK MIXTURES

### **KRENITE® S plus ESCORT XP**

KRENITE® S plus ESCORT XP may be applied for the control of Eastern red cedar and improved control of Ailanthus (tree of heaven), Ash, Cherry, Elm and Red maple.

Apply 1.5 to 3 gallons of KRENITE® S plus 1 to 2 ounces of ESCORT XP per acre. Apply a quantity of spray solution that will thoroughly and uniformly cover the target brush/trees without causing unnecessary run-off (spray to wet). If the site contains difficult-to-control species (see \*\* in "USE RATES AND PLANTS CONTROLLED" section), use the higher rates of both KRENITE® S and ESCORT XP.

Follow the use directions, precautions and restrictions on the ESCORT XP label.

### **KRENITE® S plus imazapyr**

KRENITE® S plus imazapyr herbicide (2 pounds active ingredient per gallon) may be applied for the control of Box elder, Hackberry, Persimmon, Wild pecan and Dogwood and for improved control of Ash, Black Cherry, Elm, Maple, Sassafras and Willow.

Apply 1.5 to 3 gallons of KRENITE® S plus 8 to 20 ounces of imazapyr per acre. Apply a quantity of the spray solution that will thoroughly and uniformly cover the target brush without causing unnecessary run-off (spray to wet). If the site contains difficult-to-control species (see \*\* in "USE RATES AND PLANTS CONTROLLED" section), use the higher rates of both KRENITE® S and imazapyr.

Follow the use directions, precautions and restrictions on the Imazapyr label.

### **KRENITE® S plus picloram**

KRENITE® S plus picloram (2 pound active per gallon) herbicide may be applied for the control of Hackberry, Persimmon, and Walnut for improved control of Cherry, Elm, Hickory, Locust, Oak, Poplar, Sassafras, Sumac, and Sweet gum.

Apply 1.5 to 3 gallons of KRENITE® S plus 1 to 2 pints of picloram per acre. Apply a quantity of the spray solution that will thoroughly and uniformly cover the target brush without causing unnecessary run-off (spray to wet). If the site contains difficult-to-control species (see \*\* in "USE RATES AND PLANTS CONTROLLED" section), use the higher rates of both KRENITE® S and picloram.

Follow the use directions, precautions and restrictions on the picloram label.

## SIDE TRIMMING

For control of only a portion of a plant, direct the spray solution to thoroughly cover (spray to wet) only the portion of the plant to be controlled.

Do not apply more than 6 gallons of KRENITE® S per acre when side trimming.

## CUT SURFACE APPLICATIONS

KRENITE® S may be used for controlling the re-sprouting of cut stumps of the plants listed in the "USE RATES AND PLANTS CONTROLLED" section. Control of re-sprouting in plants listed as "difficult to control" may not be as effective.

KRENITE® S may either be used undiluted or mixed with water. Use the method that is best suited for the particular application equipment. When mixing with water a ratio of no less than 1 part KRENITE® S to 1 part water on a volume basis must be used. Apply the undiluted or mixed solution to wet the area adjacent to the cambium and bark around the entire circumference and the sides of the cut stumps. The sides of the stumps should be wet down to the root collar area.

Apply with appropriate application equipment using low spray pressure. Applications can be made any time of the year, except during periods of heavy sap flow in the spring. Applications should be made soon after cutting, before the stump surface forms a layer of callous tissue (hardens off).

To prevent freezing of the spray solution, add ethylene glycol (commercial antifreeze) to the water used in preparing the spray solution. Add the antifreeze according to the manufacturer's label for preventing freezing of water at the lowest expected ambient temperature. KRENITE® S will freeze at -11°F. A 1:1 aqueous dilution of KRENITE® S will freeze at 21°F.

A spray pattern indicator may be used in the spray solution to facilitate application. The user should check the compatibility of the spray indicator with the spray solution prior to using large quantities.

## ADDITIONAL USE INSTRUCTIONS – PINE PLANTATIONS AND NON-CROP SITES

### **MIXING INSTRUCTIONS**

1. Fill spray tank 1/2 full of water.
2. With the agitator running, add the desired amount of KRENITE® S.
3. If using a tank mix partner, add the recommended amount. Follow the use precautions and directions on the tank mix partner label.
4. Add spray adjuvant as last ingredient prior to filling the spray tank with water.
5. Agitate the spray solution thoroughly.

After KRENITE® S has been thoroughly mixed in the spray tank, agitation of the spray solution is not required.

## SPRAY CLEAN-UP

Thoroughly clean all mixing and spray equipment immediately following applications of KRENITE® S. Flush tank, pump, hoses and boom with several changes of water after removing the nozzle tips and screens (clean these parts separately).

Dispose of the rinsate on a labeled site or at an approved waste disposal facility.



---

## SPRAY DRIFT MANAGEMENT

---

The interaction of many equipment- and weather-related factors determines the potential for spray drift. The applicator is responsible for considering all of these factors when making applications.

AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.

### IMPORTANCE OF DROPLET SIZE

The most effective way to reduce drift potential is to apply large droplets (>150–200 microns). The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. The presence of sensitive species nearby, the environmental conditions, and pest pressure may affect how an applicator balances drift control and coverage.

APPLYING LARGER DROPLETS REDUCES DRIFT POTENTIAL, BUT WILL NOT PREVENT DRIFT IF APPLICATIONS ARE MADE IMPROPERLY OR UNDER UNFAVORABLE ENVIRONMENTAL CONDITIONS!

See “Wind”, “Temperature and Humidity”, and “Temperature Inversions” sections of this label.

### CONTROLLING DROPLET SIZE – GENERAL TECHNIQUES

- Volume – Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.
- Pressure – Use the lower spray pressures recommended for the nozzle. Higher pressure reduces droplet size and does not improve canopy penetration. WHEN HIGHER FLOW RATES ARE NEEDED, USE A HIGHER CAPACITY NOZZLE INSTEAD OF INCREASING PRESSURE.
- Nozzle Type – Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low-drift nozzles.

### CONTROLLING DROPLET SIZE – AIRCRAFT (HELICOPTER)

- Number of Nozzles – Use the minimum number of nozzles with the highest flow rate that provide uniform coverage.
- Nozzle Orientation – Orienting nozzles so that the spray is emitted backwards, parallel to the airstream will produce larger droplets than other orientations.
- Nozzle Type – Solid stream nozzles (such as disc and core with swirl plate removed) oriented straight back produce larger droplets than other nozzle types.

### BOOM LENGTH AND HEIGHT

- Boom Length (helicopter) – For helicopters use a boom length and position that prevents droplets from entering the rotor vortices.
- Boom Height (helicopter) – Application more than 10 feet above the canopy increases the potential for spray drift.
- Boom Height (ground) – Setting the boom at the lowest height which provides uniform coverage reduces the exposure of droplets to evaporation and wind. The boom should remain level with the crop and have minimal bounce.

### WIND

Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given wind speed. AVOID GUSTY OR WINDLESS CONDITIONS.

**Note:** Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

### TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, set up equipment to produce larger droplets to reduce effects of evaporation.

### TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

---

## SHIELDED SPRAYERS

---

Shielding the boom or individual nozzles can reduce the effects of wind. However, it is the responsibility of the applicator to verify that the shields are preventing drift and not interfering with uniform deposition of the product.

### IMPORTANT PRECAUTIONS – PINE PLANTATIONS AND NON-CROP SITES

- Cutting of treated stems of brush before they are completely dead may result in sprouting.
- Do not use for the control of woody plants on lawns, walks, driveways, tennis courts or similar areas.
- Drift or spray mist contact with desirable trees, shrubs, or other plants may result in injury.
- Not registered for sale or use in California or Arizona.







## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Do not store below 10°F. Store product in original container only. Store in a cool, dry place.

**PESTICIDE DISPOSAL:** Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER HANDLING: Refer to the Net Contents section of this product's labeling for the applicable "Nonrefillable Container" or "Refillable Container" designation.**

### **Nonrefillable Rigid Plastic and Metal Containers (Capacity Equal to or Less Than 5 Gallons):**

Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

### **Nonrefillable Rigid Plastic and Metal Containers (Capacity Greater Than 5 Gallons):**

Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then, for Plastic Containers, offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances.

For Metal Containers, offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

### **Nonrefillable Rigid Plastic and Metal Containers, e.g., Intermediate Bulk Containers (IBC) [Size or Shape Too Large to be Tipped, Rolled or Turned Upside Down]:**

Nonrefillable container. Do not reuse or refill this container. Clean container promptly after emptying the contents from this container into application equipment or mix tank and before final disposal using the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top, bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration and/or spray volume. If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain, pour or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then, for Plastic Containers, offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration. For Metal Containers, offer for recycling, if available, or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

### **All Refillable Containers:**

Refillable container.

Refilling Container: Refill this container with KRENITE® S containing ammonium salt of fosamine only. Do not reuse this container for any other purpose. Cleaning before refilling is the responsibility of the refiller. Prior to refilling, inspect carefully for damage such as cracks, punctures, abrasions, worn out threads and closure devices. If damage is found, do not use container, contact Albaugh, Inc. at the number below for instructions. Check for leaks after refilling and before transporting. If leaks are found, do not reuse or transport container; contact Albaugh, Inc. at the number below for instructions.

Disposing of Container: Do not reuse this container for any other purpose other than refilling (see proceeding). Cleaning the container before final disposal is the responsibility of the person disposing of the container. To clean the container before final disposal, use the following pressure rinsing procedure. Insert a lance fitted with a suitable tank cleaning nozzle into the container and ensure that the water spray thoroughly covers the top, bottom and all sides inside the container. The nozzle manufacturer generally provides instructions for the appropriate spray pressure, spray duration and/or spray volume. If the manufacturer's instructions are not available, pressure rinse the container for at least 60 seconds using a minimum pressure of 30 PSI with a minimum rinse volume of 10% of the container volume. Drain, pour or pump rinsate into application equipment or rinsate collection system. Repeat this pressure rinsing procedure two more times. Then, for Plastic Containers, offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration. Do not burn, unless allowed by state and local ordinances. For Metal Containers, offer for recycling, if available, or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

Do not transport if container is damaged or leaking. If the container is damaged, leaking or obsolete, or in the event of a major spill, fire or other emergency, contact Albaugh, Inc. at 1-800-441-3637, day or night.





**NOTICE TO BUYER:** Purchase of this material does not confer any rights under patents of countries outside of the United States.

## LIMITATION OF WARRANTY AND LIABILITY

**NOTICE:** Read This Limitation of Warranty and Liability Before Buying or Using This Product. If the Terms Are Not Acceptable, Return the Product at Once, Unopened, and the Purchase Price Will Be Refunded.

It is impossible to eliminate all risks associated with the use of this product. Such risks arise from weather conditions, soil factors, off-target movement, unconventional fanning techniques, presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of Albaugh, Inc. These risks can cause: ineffectiveness of the product; crop injury, or; injury to non-target crops or plants.

Albaugh, Inc. does not agree to be an insurer of these risks. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, WHEN YOU BUY OR USE THIS PRODUCT, YOU AGREE TO ACCEPT THESE RISKS.**

Albaugh, Inc. warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated in the Directions for Use, subject to the inherent risks described above, when used in accordance with the Directions for Use under normal conditions.

**ALBAUGH, INC. MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.**

**TO THE FULLEST EXTENT PERMITTED BY LAW, IN NO EVENT SHALL ALBAUGH, INC. OR SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. BUYER'S OR USER'S BARGAINED-FOR EXPECTATION IS CROP PROTECTION. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE LIABILITY OF ALBAUGH, INC. OR SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, TORT OR STRICT LIABILITY), WHETHER FROM FAILURE TO PERFORM OR INJURY TO CROPS OR OTHER PLANTS, AND RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT, OR AT THE ELECTION OF ALBAUGH, INC. OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Albaugh, Inc. or its Ag Retailer must have prompt notice of any claim so that an immediate inspection of buyer's or user's growing crops can be made. Buyer and all users shall promptly notify Albaugh, Inc. or an Albaugh, Inc. Ag Retailer of any claims, whether based on contract, negligence, strict liability, other tort or otherwise or be barred from any remedy.

This Limitation of Warranty and Liability may not be amended by any oral or written agreement.

KRENITE® and AgriStar® are registered trademarks of Albaugh, Inc.  
ESCORT® is a registered trademark of E.I. DuPont de Nemours and Company.

